



DIVISION OF AGRICULTURE
RESEARCH & EXTENSION

University of Arkansas System

Introduction to Food Safety and Microbiology


Food Safety

Divided into causative categories called “hazards”

- Biological Hazards - bacteria, molds, natural occurring toxins
- Chemical Hazards – chemicals like petroleum, herbicides, pesticides, heavy metals.
- Physical Hazards – glass, rocks, wood splinters


1990 vs 1997

1900



- Tuberculosis
- Pneumonia
- Diarrhoea
- Heart disease
- Liver disease
- Injuries
- Stroke
- Cancer
- Bronchitis
- Diphtheria

1997




- Heart disease
- Cancer
- Stroke
- Chronic lung disease
- Unintentional injuries
- Pneumonia / influenza
- Diabetes
- Suicide
- Chronic kidney disease
- Chronic liver disease

Figure 2 – The ten leading causes of death in the United States in 1900 and 1997. From “Changing Patterns of Infectious Disease”, Mitchell L. Cohen, *Nature*, Volume 406, 17 August, 2002. pp 762-767.





Food Microbiology



Definition

- Microorganisms: Organisms such as bacteria, parasites, viruses, yeasts, and molds
 - Usually too small to be seen by the naked eye






Where are microorganisms?

- Soil & Water
- Plants/Products
- Utensils/Equipment
- Gastrointestinal Tract
- Food Handlers
- Animal Feeds
- Animal Hides
- Air & Dust
- EVERYWHERE!



Sources of Fecal Contamination of Food




Microorganisms in Food

Microorganisms are important in many different ways:


- Pathogenic, or disease causing, microorganisms can cause illness
- Spoilage microorganisms cause a food to smell, taste, and look unacceptable
- Fermentation microorganisms produce a desired food product
- Other microorganisms do nothing in foods

Bacterial Classifications

- Bacteria can be classified in a number of ways.
*Their shape-Round=cocci; Elongated=rods



Spores



- Sporeforming; Some rod shaped bacteria are sporeformers. This is a dormant stage in their life cycle. These spores have the ability to survive a wide range of environmental extremes. They can survive heating up to 212⁰F and are resistant to most chemicals including sanitizing solutions. The most noteworthy sporeformer is Clostridium botulinum.
- Spores ⇒ dormant state
 - Much more resistant to environmental stresses (heat, cold, chemicals)
- Vegetative state ⇒ active state
 - More susceptible to inactivation

Temperature and Growth

- PSYCHROPHILE:

Grow from 1-20°C

EXAMPLES: *Pseudomonas*, *Flavobacterium*, *Alcaligenes*

- PSYCHROTROPHIC:

Grow best at 37°C, but can grow at refrigeration (3-7°C)

EXAMPLE: *Listeria monocytogenes*

- MESOPHILE:

Optimum temperature 20-40°C


Group containing most human pathogens

EXAMPLES: *E. coli*, *Salmonella*, *Clostridium botulinum*

- THERMOPHILE:

Optimum temp >45°C

EXAMPLE: *Bacillus stearothermophilus*



Low and High Temperatures

Low Temperatures



- Refrigeration (40-45°F) slows or stops bacterial growth
- Freezing stops bacterial growth

High Temperatures

- Heating (165°F or higher) destroys bacteria for immediate service foods served in restaurants and homes.
- Thermal processing of shelf stable foods (180 - 250°F) destroys bacteria for longer shelf-life foods – temperature dependent upon product acidity
- Low acid canned foods – inactivation of *C. botulinum*


Reproduction of Bacterial Cells

- Reproduced by division
- Referred to as “growth”
- Under optimum conditions a cell divides every 20-30 minutes



Growth of Bacteria

- Binary Fission



Controlling Growth

F – Food

A – Acidity

T – Temperature

T – Time

O – Oxygen

M – Moisture




Controlling Growth



Food – Nutrients Content


- For growth, microorganisms require the following:
 - Water
 - Source of energy
 - Source of nitrogen
 - Vitamins & related growth factors
 - Minerals

Food Acidity








- Optimal pH for growth: 6.0 – 8.0
- Disease causing bacteria: 4.6 - 9.5
- Spoilage bacteria: 1.5 - 9.5

pH – Growth Range in Foods



Mild heat required, since spores are inhibited by acid.

High heat required in order to destroy spores.



Controlling Growth




T = Temperature


- Optimal Growth
 - Thermophiles – like hot conditions
 - Mesophiles – like warm conditions (around body temperature)
 - Psychrotrophs – can grow at refrigeration temperatures
- Most pathogens are mesophiles

Temperature Classifications

- *Based on optimum temperature for growth;
- *Psychro=cold
- *Meso = middle
- *Thermo= warm
- *Trophic =growing
- *Duric=withstand
- *Phil or philic-prefers or loves


Categories of Microbes Based on Temperature Range





Controlling Growth

Temperature Danger Zone



Copyright © International Association for Food Protection

Controlling Growth


- Temperature lower than 41° F
 - Bacteria cease to multiply but do not die
 - Freezing can cause cell wall damage
- Temperature higher than 135 ° F
 - Bacteria die if heated for a sufficient time
 - Increased destruction with longer times and higher temperatures

Controlling Growth





TIME

- Under optimal conditions, some bacteria can double every 20 minutes.
- The colder the storage temperature, the longer the potential shelflife.
- Potentially hazardous foods should not remain in the danger zone (50-140⁰F) for more than 4 hours during the entire food handling process.


Growth of Bacteria



Generation Time Under Optimal Conditions (at 37°C)

Organism	Generation Time (min)	
<i>Bacillus cereus</i>	28	
<i>Escherichia coli</i>	12.5	
<i>Staphylococcus aureus</i> (causes many infections: toxic shock syndrome one example)	27-30	
<i>Mycobacterium tuberculosis</i> (agent of Tuberculosis)	792 - 932	

Bacterial Growth Phases



O = Oxygen

Based on oxygen requirements;

Aerobic-Need oxygen to grow

Anaerobic-Can grow only if oxygen is
absent


Facultative-Can grow with or without
oxygen.

Controlling Growth


Oxygen

- Tolerance to oxygen in the surrounding environment


Anaerobic



Aerobic



Facultative Anaerobes




Controlling Growth

Moisture – Water Activity

- Water Activity (a_w) is the measure of “free” water available to the microorganism for growth

Water Activity

- A_w is affected by the presence of solutes (sugars and salts)




- Lowering a_w will reduce the ability for microorganisms to grow

Water Activity

Moisture in a food system that is available for microbial growth and chemical reactions – the relative humidity of a food

- 0.98 - 0.995 most foods.
- 85 - .995 for disease causing organisms range .
- 60 - .995 for spoilage organisms



WATER ACTIVITY

- Aw/ Microorganism

1.0-0.95 Bacteria

0.95-0.91 Bacteria

0.91-0.87 Yeasts

0.87-0.80 Molds

0.30-0.20 No microorganisms
 proliferate

- Foods

Meat, fish, sausage, milk


Moist cheeses, cured meat (ham),
fruit juice conc

Fermented sausages (salami), dry
cheeses, margarine

Juice conc, syrups, flour, fruit cakes,
honey, jellies, preserves


Cookies, crackers, bread crusts

Water Activity: Foods and Microbial Growth



Controlling Growth

Water Activity of Common Foods



Controlling Growth

SUMMARY:

- Microorganisms can grow on food and in the environment under the right conditions
- Remember F-A-T-T-O-M
 - Food, Acidity, Temperature, Time, Oxygen, and Moisture



Interventions

Interventions – Combined Effects

“Hurdles Concept”

Predictive microbiology
The study of interactive
effects of factors
effecting microbial
growth

- ◆ Additive effects
- ◆ Synergistic effects
- ◆ Antagonistic effects




Fig. 4.1 Hurdle effect, illustrated using four food products as example.
(From Leistner 1978.)

Food Industry Interventions

Ingredients

- Have good specifications and control of incoming ingredients

Process

- Have an adequate thermal process to destroy microorganisms
- Have an adequate packaging system to protect your product

After Processing

- Have control of distribution and a system for tracking and recalling
- Provide necessary information for consumers (labeling)

Farm to Fork Food Safety and Quality Programs

- HACCP, GMPs, Sanitation

Ingredient Control

- Specifications for ingredient make-up, quality, physical and microbial contaminant levels
 - » Letter of guarantee
 - » Certificate of analysis

Thermal Processing Control


Thermal Processing

- Cooking
- Pasteurization
- Commercial sterility (shelf-stable) processes
 - Retort systems, aseptic processing, and hot filling



Package Integrity

- Measurements are made to insure the package is of good sanitary quality and can maintain a hermetic seal (such as visual inspections and torque measurements for your containers)



Thermal Processing Control

Low acid foods (pH>4.6, Aw>.85)

- Should have a process to eliminate 1,000,000,000,000 spores of *Clostridium botulinum*
- Examples: Most meat, vegetable, and dairy products
- Process often at 230°F or above (retorts, aseptic processing)

Acidified foods (pH altered to ≤4.6, Aw>.85)

- Not required to have a process to eliminate *Clostridium botulinum*
- Examples: Pickled products, mayonnaise
- Process often 180 - 205°F (pasteurization, hot filling)

High acid foods (pH ≤4.6, Aw>.85)

- Not required to have a heat process.
- Examples: sliced oranges, condiments
- Process often 180 - 205°F (pasteurization, hot filling)

Pathogens

General Roles of Microorganisms

- Commensal
 - ubiquitous
 - harmless or beneficial
- Spoilage
 - Cause food to become inedible due to changes in color, flavor, odor, appearance or texture.
 - Grow to high levels and break down food components
 - Commensal organisms that have reached high populations (10^5 - 10^7 CFU/g)
 - Different products have different spoilage flora

General Characteristics of Microorganisms

- Beneficial
 - Used as an aid in producing desirable characteristics in food
- Pathogens
 - Cause foodborne illness
 - Microorganisms that are usually associated with the presence of pathogens are called “Indicators”. Most E coli do not cause illness.

Food Safety vs. Food Quality

- Food safety controls **HAZARDS** to the consumer.
 - A **foodborne hazard** is a biological, chemical, or physical property that may cause a food to be **unsafe** for human consumption
- Food quality controls deterioration of food to an unacceptable state

Foodborne Illness

- Foodborne illness in the United States is associated with:
 - 46 million illnesses a year
 - 325,000 hospitalizations a year
 - 3,000 deaths a year
 - A loss of \$10-83 billion in pain & suffering, reduced productivity, and medical costs

Common Foodborne Pathogens

Bacteria

- *E. coli* 0157:H7
- *Salmonella* spp.
- *Staphylococcus aureus*
- *Listeria monocytogenes*
- *Campylobacter jejuni*
- *Shigella* spp.

Viruses

- Norovirus
- Rotovirus
- Hepatitis A

Parasites

- ◆ *Cryptosporidium parvum*
- ◆ *Giardia lamblia*
- ◆ *Cyclospora*

80-90% of Foodborne Illnesses from Bacteria come from just 4 Bacteria

- Campylobacter
- Salmonella
- Clostridium perfringens
- Staphylococcus aureus

Illness Mechanisms

- Infection
 - Microorganisms are ingested and then cause illness
- Intoxication
 - Toxins are produced by the pathogen, usually in the food. When food is consumed, illness occurs.
 - Even if microorganisms are killed, toxin can still remain the food

Bacterial Pathogens of Concern

- *E. coli* O157:H7
- *Salmonella*
- *Listeria*
- *Campylobacter*
- *Staphylococcus aureus*
- *Clostridium botulinum*


Which Bacteria are Responsible?

Pathogen	Cases	Deaths
<i>Campylobacter jejuni</i>	4,000,000	200-1000
<i>Salmonella</i>	2,000,000	500-2000
<i>Stapylococcus aureas</i>	1,500,000	1200
<i>Escherichia coli</i> O157:H7	725,000	100-200
<i>Clostridium</i> spp.	10,000	100
<i>Listeria monocytogenes</i>	1500	250-500

E. coli O157:H7

- Hemorrhagic colitis
- Cause: infection
- Incubation: 2-4 days
- Symptoms: diarrhea (blood), HUS, TPP
- Contaminant: milk, meat, fruits, vegetables, water





Salmonella


- Salmonellosis
- >2000 strains, 10 = foodborne illness
- Cause: infection
- Incubation: 6-48 hours
- Symptoms: nausea, fever, diarrhea, arthritis
- Contaminant: milk, meat, eggs

Listeria monocytogenes

- Listeriosis
- Cause: infection
- Incubation: 2 days - 3 weeks
- Symptoms: vomiting, diarrhea
 - meningitis, septicemia, miscarriage
- Contaminant: vegetables, milk, cheese, meat, seafood




Campylobacter jejuni




- Campylobacteriosis
- Cause: infection
- Incubation: 2 - 5 days
- Symptoms: nausea, fever, diarrhea (blood)
- Contaminant: milk, meat, water

Staphylococcus aureus



- Staphyloenterotoxigenesis
- Cause: intoxication
 - (1 mg toxin = 100,000 cfu/g)
- Incubation: 1-6 hours
- Symptoms: nausea, fever, diarrhea
- Contaminant: milk, meat, eggs

Clostridium botulinum



- Botulism
- Cause: intoxication (spores - neurotoxin)
- Incubation: 18 -36 hours
- Symptoms: weakness, vertigo
 - difficulty in speaking, swallowing, breathing
- Contaminant: pH >4.6, low oxygen foods

Prevention of Foodborne Illness

- 1) Cook- Cook all meat, poultry and eggs to at least 160F. Other than spore-forming bacteria, all bacteria, parasites and viruses are killed quite easily with heating to 160F.
- 2) Avoid Cross-Contamination- Do not cross-contaminate one food with another. Keep raw food totally separated from cooked product. Clean utensils and work areas etc in between working raw and cooked product. Constantly be thinking of how microorganisms get from raw to cooked products.

Prevention of Foodborne Illnesses

- 3) Chill Foods- Keep foods cold. After cooking, chill foods as rapidly as possible. Remember that cooking has destroyed most of the bacteria but spore formers, that are resistant to cooking may become very active and can proliferate rapidly.
- 4) Cleaning-Wash fruits and vegetables and all foods possible. In addition, continually wash work areas. Use only treated or tested water.

Prevention of Foodborne Illnesses

5) Personal Hygiene- People working with foods should wash their hands regularly, wear hairnets, plastic gloves etc. In addition, food handlers should not work with food if they have a boil, open sores or feel sick themselves

Spoilage Organisms

- Bacterial (hundreds of bacteria cause spoilage)
 - *Erwinia*, *Pseudomonas*, *Flavobacteria*, & *Enterobacter* spp.
 - Lactic acid bacteria
- Fungal
 - *Penicillium*, *Aspergillus*, *Fusarium*, and *Candida*

CONCLUSIONS

- Food Microbiology is huge area
- FATTOM
- Pathogens – Food Safety
- Spoilage microorganisms – costly